

Some properties of subspaces-hypercyclic operators

Nareen Sabih ^{*1} and Adem Kılıçman ^{†2}

^{1,2}Department of Mathematics, University Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

Abstract

In this paper, we answer a question posed in the introduction of [10] positively, i.e, we show that if T is $\mathcal{M}$ -hypercyclic operator with $\mathcal{M}$ -hypercyclic vector x in a Hilbert space $\mathcal{H}$, then $P(\text{Orb}(T, x))$ is dense in the subspace $\mathcal{M}$ where P is the orthogonal projection onto $\mathcal{M}$. Furthermore, we give some relations between $\mathcal{M}^\perp$ -hypercyclicity and the orthogonal projection onto $\mathcal{M}^\perp$. We also give sufficient conditions for a bilateral weighted shift operators on a Hilbert space $\ell^2(\mathbb{Z})$ to be subspace-hypercyclic, cosequently, there exists an operator T such that both T and T^* are subspace-hypercyclic operators. Finally, we give an $\mathcal{M}$ -hypercyclic criterion for an operator T in terms of its eigenvalues.

1 Introduction

A bounded linear operator T on a separable Hilbert space $\mathcal{H}$ is hypercyclic if there is a vector $x \in \mathcal{H}$ such that $\text{Orb}(T, x) = \{T^n x : n \geq 0\}$ is dense in $\mathcal{H}$, such a vector x is called hypercyclic for T . The first example of a hypercyclic operator on a Banach space was constructed by Rolewicz in 1969 [13]. He showed that if B is the backward shift on $\ell^p(\mathbb{N})$ then λB is hypercyclic if and only if $|\lambda| > 1$. The hypercyclicity concept was probably born with the thesis of Kitai in 1982 [8] who introduced the hypercyclic criterion to ensure the existence of hypercyclic operators. For more information on hypercyclic operators we refer the reader to [1, 5]

In 2011, Madore and Martínez-Avendaño [10] studied the density of the orbit in a non-trivial subspace instead of the whole space, this phenomenon is called the subspace-hypercyclicity. For the series of references on subspaces-hypercyclic operators see [7, 9, 10, 12]

Definition 1.1. [10] Let $T \in \mathcal{B}(\mathcal{H})$ and $\mathcal{M}$ be a closed subspace of $\mathcal{H}$. Then T is called $\mathcal{M}$ -hypercyclic or subspace-hypercyclic operator for a subspace $\mathcal{M}$ if there exists a vector $x \in \mathcal{H}$ such that $\text{Orb}(T, x) \cap \mathcal{M}$ is dense in $\mathcal{M}$. We call x an $\mathcal{M}$ -hypercyclic vector for T .

Definition 1.2. [10] Let $T \in \mathcal{B}(\mathcal{H})$ and $\mathcal{M}$ be a closed subspace of $\mathcal{H}$. Then T is called $\mathcal{M}$ -transitive or subspace-transitive for a subspace $\mathcal{M}$ if for each pair of non-empty open sets U_1, U_2 of $\mathcal{M}$ there exists an $n \in \mathbb{N}$ such that $T^{-n}U_1 \cap U_2$ contains a non-empty relatively open set in $\mathcal{M}$.

Theorem 1.3. [10] Every $\mathcal{M}$ -transitive operator on $\mathcal{H}$ is $\mathcal{M}$ -hypercyclic.

Theorem 1.4. [11] Every hypercyclic operator is subspace-hypercyclic for a subspace $\mathcal{M}$ of $\mathcal{H}$.

^{*}nareen_bamerni@yahoo.com

[†]akilicman@yahoo.com

In the present paper, we answer positively a question posed by Madore and Martínez-Avendaño in the introduction of [10]. In particular, Theorem 2.1 shows that if P is the orthogonal projection onto a subspace $\mathcal{M}$, then $P(\text{Orb}(T, x))$ will be dense in $\mathcal{M}$ whenever T is $\mathcal{M}$ -hypercyclic. Furthermore, we give some relations between $\mathcal{M}^\perp$ -hypercyclicity and the orthogonal projection onto $\mathcal{M}^\perp$.

Through Theorem 2.4, we give a set of sufficient conditions for a bilateral weighted shift on a Hilbert space $\ell^2(\mathbb{Z})$ to be $\mathcal{M}$ -hypercyclic operator. Like the hypercyclicity case, Example 2.7 shows that there is an operator T such that both T and its adjoint are subspace-hypercyclic; however, we don't know whether they are subspace-hypercyclic for the same subspace or not. According to which an operator having a large supply of eigenvectors is hypercyclic, Godefroy-Shapiro [4] exhibited a hypercyclic criterion which is called ‘‘Godefroy-Shapiro Criterion’’. We extend such a criterion to a subspaces and we call it ‘‘Spectrum $\mathcal{M}$ -hypercyclic Criterion’’. We give the Example 2.9 to show that the $\mathcal{M}$ -hypercyclic criterion is a stronger result than Spectrum $\mathcal{M}$ -Hypercyclic Criterion.

2 Main results

The set of all $\mathcal{M}$ -hypercyclic operators is denoted by $HC(\mathcal{H}, \mathcal{M})$ and the set of all $\mathcal{M}$ -hypercyclic vectors of T is denoted by $HC(T, \mathcal{M})$.

To prove the following results, the reader should be convenient with the properties of the projection map P , see [2].

In the introduction of [10], the authors asks the possibility of $P(\text{Orb}(T, x))$ to be dense in $\mathcal{M}$ where P is the orthogonal projection onto $\mathcal{M}$.

Theorem 2.1. *If $x \in HC(T, \mathcal{M})$ and $P : \mathcal{H} \rightarrow \mathcal{M}$ is the orthogonal projection onto $\mathcal{M}$, then $P(\text{Orb}(T, x)) \cap \mathcal{M}$ is dense in $\mathcal{M}$.*

Proof. Since $x \in HC(T, \mathcal{M})$, then there exist a sequence $\{n_k\}$ of positive numbers such that $\text{Orb}(T, x) \cap \mathcal{M} = \{T^{n_k}x : k \in \mathbb{N}\}$ is dense in $\mathcal{M}$. Since

$$\{T^{n_k}x : k \in \mathbb{N}\} \subseteq P(\text{Orb}(T, x)) \cap \mathcal{M},$$

then $P(\text{Orb}(T, x)) \cap \mathcal{M}$ is dense in $\mathcal{M}$. □

The following examples gives an application to Theorem 2.1

Example 2.2. *Let B be the backward shift on $\ell^2(\mathbb{N})$ and $\mathcal{M}$ be the subspace of $\ell^2(\mathbb{N})$ consisting of all sequences with zeroes on the even entries; that is*

$$\mathcal{M} = \{\{x_n\}_{n=0}^\infty \in \ell^2(\mathbb{N}) : x_{2k} = 0 \text{ for all } k\}$$

Since $2B \in HC(\mathcal{M}, \mathcal{H})$ see [10, Example 3.8.], then there exists an $x \in HC(2B, \mathcal{M})$. We may, and will, assume that $x = (a_0, 0, a_1, 0, \dots) \in \mathcal{M}$, then for all $n \geq 0$,

$$(2B)^{2n+1}x = 2^{2n+1}(0, a_{2n+2}, 0, a_{2n+4}, \dots)$$

and

$$(2B)^{2n}x = 2^{2n}(a_{2n}, 0, a_{2n+2}, 0, a_{2n+4}, \dots).$$

Therefore

$$\text{Orb}(2B, x) \cap \mathcal{M} = (2B)^{2n}x \text{ is dense in } \mathcal{M} \tag{1}$$

Moreover

$$P(\text{Orb}(T, x)) = \begin{cases} 0 & \text{if } n \text{ is odd,} \\ (2B)^n x & \text{if } n \text{ is even.} \end{cases}$$

Thus, it is clear from eq. (1) that $(2B)^{2n}x \subset P(\text{Orb}(T, x)) \cap \mathcal{M} = (2B)^{2n}x \cup \{0\}$ for all $n \geq 0$ and this gives the proof.

Proposition 2.3. Let $\mathcal{M}^\perp$ be an invariant subspace under T , P be the orthogonal projection onto $\mathcal{M}^\perp$ and $T \in HC(\mathcal{H}, \mathcal{M}^\perp)$, then

1. $S \in HC(\mathcal{H}, \mathcal{M}^\perp)$ where $S = PT : \mathcal{M}^\perp \rightarrow \mathcal{M}^\perp$,
2. $\bar{T} \in HC(\mathcal{H}/\mathcal{M})$ where $\bar{T} : \mathcal{H}/\mathcal{M} \rightarrow \mathcal{H}/\mathcal{M}$.

Proof. (1): Let $x \in HC(T, \mathcal{M}^\perp)$ where $x = u + v$; $u \in \mathcal{M}$ and $v \in \mathcal{M}^\perp$, then $Px = v$. Without loss of generality, we may assume that $x \in \mathcal{M}^\perp$, then

$$\begin{aligned} \text{Orb}(PT, v) \cap \mathcal{M}^\perp &= \{v, PTv, (PT)^2v, \dots\} \cap \mathcal{M}^\perp \\ &= \{v, PTv, PT^2v, \dots\} \cap \mathcal{M}^\perp \\ &= \{Px, PTPx, PT^2Px, \dots\} \cap \mathcal{M}^\perp \\ &= \{Px, PTx, PT^2x, \dots\} \cap \mathcal{M}^\perp \\ &= P(\text{Orb}(T, x)) \cap \mathcal{M}^\perp \\ &= P(\text{Orb}(T, x)) \end{aligned}$$

Since P is continuous, then

$$\begin{aligned} \overline{(\text{Orb}(PT, v) \cap \mathcal{M}^\perp)} &= \overline{P(\text{Orb}(T, x))} \\ &\supseteq P(\overline{\text{Orb}(T, x)}) \\ &\supseteq \mathcal{M}^\perp. \end{aligned}$$

Thus $PT \in HC(\mathcal{H}, \mathcal{M}^\perp)$.

(2): Without loss of generality, we may assume that $x \in \mathcal{M}^\perp$. By part (1), $PT \in HC(\mathcal{H}, \mathcal{M}^\perp)$ where $PT : \mathcal{M}^\perp \rightarrow \mathcal{M}^\perp$ and since $(PT)^n x = PT^n x$ for all $n \geq 0$, then $P(\text{Orb}(T, x))$ is dense in $\mathcal{M}^\perp$. Now, we will show that $x + \mathcal{M} \in HC(\bar{T})$,

$$\begin{aligned} \overline{\{(\bar{T}^n(x + \mathcal{M})) : n \geq 0\}} &= \overline{\{(T^n x + \mathcal{M}) : n \geq 0\}} \\ &= \overline{\{PT^n x : n \geq 0\}} + \mathcal{M}, \text{ since } T^n x \in \mathcal{M}^\perp \\ &= \overline{P(\text{Orb}(T, x))} + \mathcal{M} \\ &= \mathcal{M}^\perp / \mathcal{M} = \mathcal{H}/\mathcal{M}. \end{aligned}$$

□

The next Theorem gives sufficient conditions for a bilateral weighted shift operator on the Hilbert space $\ell^2(\mathbb{Z})$ to satisfy the $\mathcal{M}$ -hypercyclic criterion. We will suppose that

$$\mathcal{B} = \{e_{m_r} : r \in \mathbb{N}\}$$

is a Schauder basis for $\mathcal{M}$, where $m_r \in \mathbb{Z}$. Let T be the bilateral forward weighted shift operator with a weigh sequence w_n , then $T(e_{m_r}) = w_{m_r} e_{m_r+1}$ for all $r \in \mathbb{N}$. We may define a right inverse (backward shift) S to T as follows: $S(e_{m_r}) = \frac{1}{w_{m_r-1}} e_{m_r-1}$. Observe that $TSe_{m_r} = e_{m_r}$ for all $r \in \mathbb{N}$. If T is invertable then $T^{-1} = S$. Also note that for all $r \in \mathbb{N}$ and $k \geq 0$, we have

$$T^k(e_{m_r}) = \left(\prod_{j=m_r}^{m_r+k-1} w_j \right) e_{m_r+k} \text{ and } S^k(e_{m_r}) = \left(\prod_{j=m_r-1}^{m_r-k} \frac{1}{w_j} \right) e_{m_r-k}$$

First we need the following lemma.

Lemma 2.4. *Suppose that T is an invertible bilateral weighted shift, $\{n_k\}$ is a sequence of positive integers, $n_k \rightarrow \infty$ and $\mathcal{M}$ is closed subspace such that $T^{n_k}\mathcal{M} \subseteq \mathcal{M}$. If there exists an $i \in \mathbb{N}$ such that $T^{n_k}e_{m_i} \rightarrow 0$ as $k \rightarrow \infty$, then $T^{n_k}e_{m_r} \rightarrow 0$ for all $r \in \mathbb{N}$.*

Proof. Since $T^{n_k}\mathcal{M} \subseteq \mathcal{M}$, the proof is similar to the proof of [3, Lemma 3.1]. $\square$

Theorem 2.5. *Let T be an invertible bilateral forward weighted shift in the Hilbert space $\ell^2(\mathbb{Z})$ with a positive weight sequence $\{w_n\}_{n \in \mathbb{Z}}$ and $\mathcal{M}$ be a subspace of $\ell^2(\mathbb{Z})$, then T satisfies the $\mathcal{M}$ -hypercyclic criterion if there exists a sequence of positive integers $\{n_k\}$ such that $T^{n_k}\mathcal{M} \subseteq \mathcal{M}$ and $i \in \mathbb{N}$ such that*

$$\lim_{k \rightarrow \infty} \prod_{j=m_i}^{m_i+n_k-1} w_j = 0 \text{ and } \lim_{k \rightarrow \infty} \prod_{j=m_i-1}^{m_i-n_k} \frac{1}{w_j} = 0$$

Proof. We will verify the $\mathcal{M}$ -hypercyclic criterion with $D = D_1 = D_2 =$ be a dense subset of $\mathcal{M}$ consisting of all finite sequences, that is, those sequences only have finite number of nonzero entries. Let $x \in D$, then it suffices to show that $x = e_{m_r}$ for $r \in \mathbb{N}$. Furthermore, by Lemma 2.4 it suffices to show that $T^{n_k}e_{m_i} \rightarrow 0$ and $S^{n_k}e_{m_i} \rightarrow 0$. But that is clear because

$$\|T^{n_k}e_{m_i}\| = \prod_{j=m_i}^{m_i+n_k-1} w_j \rightarrow 0 \text{ and } \|S^{n_k}e_{m_i}\| = \prod_{j=m_i-1}^{m_i-n_k} \frac{1}{w_j} \rightarrow 0. \text{ Moreover it is clear that } T^{n_k}S^{n_k}x = x. \text{ By taking } x_k = S^{n_k}, \text{ it is clear that the conditions of } \mathcal{M}\text{-hypercyclic criterion are satisfied. } \square$$

By the same way we can characterize the $\mathcal{M}$ -hypercyclic backward weighted shifts since they are unitarily equivalent to forward shifts.

Corollary 2.6. *If S is an invertible bilateral backward weighted shift in the Hilbert space $\ell^2(\mathbb{Z})$ with a positive weight sequence $\{w_n\}_{n \in \mathbb{Z}}$ and $\mathcal{M}$ is a subspace of $\ell^2(\mathbb{Z})$, then T satisfies the $\mathcal{M}$ -hypercyclic criterion if there exists a sequence of positive integers $\{n_k\}$ such that $T^{n_k}\mathcal{M} \subseteq \mathcal{M}$ and $i \in \mathbb{N}$ such that*

$$\lim_{k \rightarrow \infty} \prod_{j=m_i}^{m_i-n_k+1} w_j = 0 \text{ and } \lim_{k \rightarrow \infty} \prod_{j=m_i+1}^{m_i+n_k} \frac{1}{w_j} = 0$$

From the above theorem, one can construct a weight sequence $\{w_n\}$ such that $\{w_n\}$ satisfy the conditions of Theorem 2.5 and Corollary 2.6. It immediately follows that both T and T^* are subspaces-hypercyclic operators. In other words, we can show that the Herrero question [6] holds true even on a subspace of a Hilbert space.

Example 2.7. *Let T be a bilateral shift defined as in the example of Salas [14]. Then, both T and T^* are $\mathcal{M}$ -hypercyclic operators.*

Proof. The proof follows directly from theorem 1.4. In particular, there exist two subspaces $\mathcal{M}_1$ and $\mathcal{M}_2$ such that T is $\mathcal{M}_1$ -hypercyclic and T^* is $\mathcal{M}_2$ -hypercyclic. $\square$

Question 1. *If T is $\mathcal{M}_1$ -hypercyclic and T^* is $\mathcal{M}_2$ -hypercyclic. What is the relation between $\mathcal{M}_1$ and $\mathcal{M}_2$?*

Conjecture 1. *If T is $\mathcal{M}_1$ -transitive and T^* is $\mathcal{M}_2$ -transitive, then $\mathcal{M}_2 = \mathcal{M}_1^\perp$.*

Theorem 2.8 (Spectrum $\mathcal{M}$ -Hypercyclic Criterion). *Let $T \in \mathcal{B}(\mathcal{H})$ and $\mathcal{M}$ be a subspace of $\mathcal{H}$. If there is a positive integers p such that the subspace $X = \text{span} \{x \in \mathcal{H}; T^p x = \lambda x, \lambda \in \mathbb{C}, |\lambda| < 1\} \cap \mathcal{M}$, $Y = \text{span} \{x \in \mathcal{H}; T^p x = \lambda x, \lambda \in \mathbb{C}, |\lambda| > 1\} \cap \mathcal{M}$ are dense in $\mathcal{M}$. Then T is $\mathcal{M}$ -transitive.*

Proof. Let U, V be nonempty open subsets of $\mathcal{M}$. By hypothesis there exist $x \in X \cap U$ and $y \in Y \cap V$ such that

$$x = \sum_{k=1}^m a_k x_k \quad \text{and} \quad y = \sum_{k=1}^m b_k y_k$$

where $x_k \in X$ and $y_k \in Y$ which mean that $T^p x_k = \lambda_k x_k$, $|\lambda_k| < 1$ and $T^p y_k = \mu_k y_k$, $|\mu_k| > 1$, for some $a_k, b_k, \lambda_k, \mu_k \in \mathbb{C}$, $k = 1, \dots, m$. Let $z_n = \sum_{k=1}^m b_k \frac{1}{\mu_k^n} y_k$. Since as $(n) \rightarrow \infty$,

$$T^{n+p} x = \sum_{k=1}^m a_k \lambda_k^n x_k \rightarrow 0 \quad \text{and} \quad z_n \rightarrow 0$$

and since $T^{n+p} z_n = y$ for all $n \geq 0$, then there exists $N \in \mathbb{N}$ such that, for all $n \geq N$,

$$x + z_n \in U \quad \text{and} \quad T^{n+p}(x + z_n) = T^{n+p} x + y \in V.$$

It follows that for all $n \geq N$,

$$T^{n+p} U \cap V \neq \emptyset$$

Now, let $w \in \mathcal{M}$, since X is dense in $\mathcal{M}$, then there exists a sequence $(x_k) \in X$ such that $(x_k) \rightarrow w$. since $T \in \mathcal{B}(\mathcal{H})$, then $T^{n+p}(x_k) = (\lambda_k^n x_k) \rightarrow T^{n+p} w$ for all $n \geq 0$. Since $|\lambda_k| < 1$ for all $k \geq 0$, then $(\lambda_k^n x_k) \rightarrow 0$ as $n \rightarrow \infty$. Thus, $0 = T^{n+p} w \in \mathcal{M}$ and we get $T^{n+p}(\mathcal{M}) \subseteq \mathcal{M}$ for all $n \geq N$. This show that T is $\mathcal{M}$ -transitive. $\square$

We give an example showing that the $\mathcal{M}$ -hypercyclic criterion is a stronger result than Spectrum $\mathcal{M}$ -Hypercyclic Criterion.

Example 2.9. *Consider the bilateral weighted forward shift $T = F : \ell^2(\mathbb{Z}) \rightarrow \ell^2(\mathbb{Z})$, with the weight sequence*

$$w_n = \begin{cases} \frac{1}{2} & \text{if } n \geq 0, \\ 3 & \text{if } n < 0. \end{cases}$$

Let $\mathcal{M}$ be the subspace of $\ell^2(\mathbb{Z})$ consisting of all sequences with zeroes on the even entries; that is,

$$\mathcal{M} = \{ \{a_n\}_{n=-\infty}^{\infty} \in \ell^2(\mathbb{Z}) : a_{2n} = 0, n \in \mathbb{Z} \}$$

then T satisfies $\mathcal{M}$ -hypercyclic criterion, but not spectrum $\mathcal{M}$ -hypercyclic criterion.

Proof. Applying Theorem 2.5 with $n_k = 2k$ and $m_i = 1$, we get

$$\lim_{k \rightarrow \infty} \prod_{j=1}^{2k} w_j = \lim_{k \rightarrow \infty} \prod_{j=1}^{2k} \frac{1}{2} = 0 \quad \text{and} \quad \lim_{k \rightarrow \infty} \prod_{j=-1}^{-2k} \frac{1}{w_j} = \lim_{k \rightarrow \infty} \prod_{j=1}^{2k} \frac{1}{3} = 0$$

It can be easily deduced from the definition of $\mathcal{M}$ that for each sequence $x \in \mathcal{M}$ and each $k \in \mathbb{Z}$, the sequence $T^{2k} x$ will have a zero entry on all even positions. It follows that $T^{n_k} \mathcal{M} \subseteq \mathcal{M}$. Thus T satisfies the $\mathcal{M}$ -hypercyclic criterion.

On the other hand, let $x = (\dots, 0, x_{-3}, 0, x_{-1}, 0, x_1, 0, x_3, 0, \dots)$ be a non-zero element in $\mathcal{M}$. Towards a contradiction, assume that there exist $p \in \mathbb{N}$ such that $T^p x = \lambda x$, then if p is odd

number, then we get $x = 0$; otherwise, without loss of generality we will assume that $p = 2$. The equality $T^2x = \lambda x$ implies that

$$\begin{aligned} x &= \left(\dots, \left(\frac{\lambda}{3^2} \right)^2 x_{-1}, 0, \left(\frac{\lambda}{3^2} \right) x_{-1}, 0, x_{-1}, 0, 3 \frac{1}{2\lambda} x_{-1}, 0, 3 \frac{1}{2^3 \lambda^2} x_{-1}, 0, 3 \frac{1}{2^5 \lambda^3} x_{-1}, 0, \dots \right) \\ &= \left(\dots, \left(\frac{\lambda}{3^2} \right)^2 x_{-1}, 0, \left(\frac{\lambda}{3^2} \right) x_{-1}, 0, x_{-1}, 0, 6 \left(\frac{1}{2^2 \lambda} \right) x_{-1}, 0, 6 \left(\frac{1}{2^2 \lambda} \right)^2 x_{-1}, 0, 6 \left(\frac{1}{2^2 \lambda} \right)^3 x_{-1}, 0, \dots \right) \end{aligned}$$

where $\lambda \neq 0$ and $x_{-1} \neq 0$. But then we have that

$$\|x\| = |x_{-1}| \left(1 + \sum_{n \in \mathbb{N}} \left(\frac{|\lambda|}{3^2} \right)^n + 6 \sum_{n \in \mathbb{N}} \left(\frac{1}{2^2 |\lambda|} \right)^n \right) = \infty$$

whatever the value of λ . Thus T^2 has no eigenvalues and therefore it does not satisfy the spectrum $\mathcal{M}$ -hypercyclic criterion. $\square$

References

- [1] F. Bayart, É. Matheron, *Dynamics of Linear Operators*, Cambridge University Press, 2009.
- [2] J. B. Conway. *A Course in Functional Analysis*, volume 96 of Graduate Texts in Mathematics. Springer-Verlag, 1990.
- [3] N. Feldman, *Hypercyclicity and supercyclicity for invertible bilateral weighted shifts*, Proc. Amer. Math. Soc. **131** (2003) 479-485.
- [4] G. Godefroy and J.H. Shapiro, *Operators with dense, invariant, cyclic vector manifolds*, J. Funct. Anal. 98 (1991), 229-269.
- [5] K.G. Grosse-Erdmann, A. Peris, *Linear Chaos*, Universitext, Springer, 2011.
- [6] D. A. Herrero. *Limits of hypercyclic and supercyclic operators*, J. Funct. Anal., **99** (1991) 179-190.
- [7] R.R. Jiménez-Munguía, R. A. Martínez-Avendaño, A. Peris, *Some questions about subspace-hypercyclic operators*, J. Math. Anal. Appl. 408 (2013), 209-212.
- [8] C. Kitai, *Invariant Closed Sets for Linear Operators*, Thesis, University of Toronto, 1982.
- [9] C.M. Le, *On subspace-hypercyclic operators*, Proc. Amer. Math. Soc., B9(2011), 2847-2852.
- [10] B.F. Madore, R. A. Martínez-Avendaño, *Subspace-hypercyclicity*, J. Math. Anal. Appl. 373 (2011), 502-511.
- [11] M.S. Nareen and Adem Kılıçman, *Some characterizations of the subspaces-diskcyclicity*, Submitted for publication.
- [12] H. Rezaei, *Notes on subspace-hypercyclic operators*, J. Math. Anal. Appl. 397 (2013), 428-433.
- [13] S. Rolewicz, *On orbits of elements*, Studia Math. **32** (1969), 17-22.
- [14] H. N. Salas. A hypercyclic operator whose adjoint is also hypercyclic. Proc. Amer. Math. Soc., 112 (3): 765-770, 1991.